

C57BL/6-GFP Mouse Primary Dermal Microvascular Endothelial Cells

Catalog No. mGFP-6064

Suggested Medium

M1168 Complete Endothelial Cell Medium w/ Kit – 500 ml

Product Description

C57BL/6-GFP Mouse Primary Dermal Microvascular Endothelial Cells from Cell Biologics are isolated from C57BL/6-Tg (CAG-EGFP) 1Osb/J mouse skin tissues of pathogen-free laboratory mice. Cells at passage 3 are harvested from flasks and each vial contains 0.5×10^6 cells per ml. The cells are characterized by use of fluorescence-labeled acetylated low-density lipoprotein (DiI-Ac-LDL) uptake (Catalog No. L-35353, Invitrogen), a functional marker for endothelial cells. These cells are negative for bacteria, yeast, fungi, and mycoplasma and can be expanded for 3-5 passages at a split ratio of 1:2 under the cell culture conditions specified by Cell Biologics. Repeated freezing and thawing of cells are not recommended.

Storage

Cryopreserved cells are shipped with dry ice overnight. Upon arrival, transfer frozen cells to liquid nitrogen (-180°C) immediately until ready for use. Live cell shipment is also available on request. Primary cells can never be kept at -20°C or -80°C freezer.

Authorized Uses of Cell Biologics' Products

The Cells from Cell Biologics are distributed for research purposes only. Our products are not authorized for human use, for in vitro diagnostic or therapeutic procedures. Transfer or resale of any Cell Biologics' cells or products from the purchaser to other markets, organizations or individuals is prohibited by Cell Biologics without the company's written consent. Cell Biologics' Terms and Conditions must be accepted before submitting an order.

Disclaimer

Investigators should handle the cells with caution and treat all animal cells as potential pathogens, since no test procedure can completely guarantee the absence of infectious agents.

Warranty and Liability

Cell Biologics' guarantee applies only to your purchase of Cell Biologics' Cells with Cell Biologics' Media and Coating Solution for appropriate cell culture and cell testing following Cell Biologics' online protocols within 35 days from the date of product delivery.